

67760-110

11/15/2012

1/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

Carrie M. Tackema
Regulatory Affairs Manager
Cheminova, Inc.
P.O. Box 110566
Research Triangle Park, NC 27709

NOV 15 2012

Product Name: KOVERALL™ Fungicide
EPA Reg. No.: 67760-110
Subject: Your amendment dated October 10, 2012
OPP Decision Number: 471675

Dear Ms. Tackema:

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended is acceptable provided the following changes are made:

1. On page 1, change "THIS SUPPLEMENTAL LABEL EXPIRES ON XXXXX.... AFTER THIS DATE." to "THIS SUPPLEMENTAL LABEL EXPIRES ON MARCH 31, 2014....AFTER THIS DATE."
2. On page 1, add the following immediately below the header **Directions for Use**: "It is a violation of Federal law to use this product in a manner inconsistent with its labeling."
3. On page 1 in the second line below the header **IMPORTANT**, change "...EPA-REGISTERED LABEL." to "...EPA-REGISTERED PACKAGE LABEL."
4. On page 1 in the second line of the second paragraph below the header **IMPORTANT**, change "...the Section 3 Federal product label...." to "...the container label...."

One copy of the supplemental label stamped "Accepted with comments" is enclosed for your records. Please submit one copy of the final printed label that incorporates the required changes before the product is released for shipment. If you have any questions, please contact Robert Westin by phone at (703) 305-5721 or via email at westin.robert@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mary L. Waller".

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure

SUPPLEMENTAL LABELING
ACCEPTED
with COMMENTS
In EPA Letter Dated:
11/15/2012

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. 67760-110



KOVERALL™ Fungicide

EPA Reg. No. 67760-110

THIS SUPPLEMENTAL LABEL EXPIRES ON XXXXX
AND MUST NOT BE USED OR DISTRIBUTED AFTER THIS DATE

Directions for Use

KOVERALL Fungicide is a broad-spectrum fungicide. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. The addition of an agricultural surfactant will enhance fungicide performance by providing a more uniform spray deposit, increased foliar redistribution, and enhanced fungicide retention during periods of wet weather.

IMPORTANT

BEFORE USING KOVERALL, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the WARRANTY DISCLAIMER, INHERENT RISKS OF USE, and LIMITATION OF REMEDIES on the Section 3 Federal product label.

This labeling must be in the possession of the user at the time of pesticide application.

Cole Crop (Brassica) Leafy Vegetables

Crop	Diseases Controlled	KOVERALL™ Rate per Application (lb/acre)	Application Directions	Restrictions
Broccoli and Cabbage	Alternaria Leaf Spot Downy Mildew	1.6-2.1	In plant beds or direct-seeded fields, apply 7 to 10 days after planting or earlier if disease is present. In field applications, apply as soon as disease is present and reapply as needed on a 7 to 10 day spray schedule.	Do not apply more than 12.8 lb. KOVERALL (9.6 lb ai) per acre. Preharvest Interval: Do not apply within 7 days of harvest. Do not apply this product with a U-boom device. Minimum Retreatment Interval: 7 days

Fruiting Vegetables (Except Cucurbits)

Crop	Diseases Controlled	KOVERALL Rate per Application (lb/acre)	Application Directions	Restrictions
Peppers	Anthracnose Early Blight Phomopsis Blight or Fruit Rot Alternaria Leaf Spot Downy Mildew Cercospora Leaf Spot (Frogeye Spot) Phytophthora Blight Ripe Rot	2 (west of the Mississippi River) 3 (east of the Mississippi River)	Begin application when disease appears and reapply on a 7 to 10 day spray schedule.	East of the Mississippi River, do not apply more than 19.2 lb KOVERALL (14.4 lb ai) per acre per crop and do not apply within 7 days of harvest. West of the Mississippi River, do not apply more than 12.8 lb KOVERALL (9.6 lb ai) per acre per crop and do not apply within 7 days of harvest. Do not apply this product with a U-boom device. Minimum retreatment interval: 7 days

**Lettuce
(Head and Leaf)**

Crop	Diseases Controlled	KOVERALL™ Rate per Application (lb/acre)	Application Directions	Restrictions
Lettuce	Downy Mildew Anthracnose Alternaria Leaf Spot (not for use on this pest in California)*	2	Begin applications when disease appears and reapply on a 7 to 10 day treatment schedule.	In California , do not apply more than 8.5 lb KOVERALL (6.4 lb ai) per acre per crop and do not apply within 14 days of harvest. In states other than California , do not apply more than 12.8 lb KOVERALL (9.6 lb ai) per acre per crop and do not apply within 10 days of harvest. Do not apply this product with a U-boom device. Minimum retreatment interval: 7 days

Tree Nuts

Crop	Diseases Controlled	KOVERALL™ Rate per Application (lb/acre)	Remarks (Also Refer to Directions for Use)	Restrictions
Almond	Blossom Blight (<i>Monilinia</i> spp) Shothole (<i>Stigmina</i> spp)	6.4 lb/A	Begin application at dormant to popcorn stage, full bloom or petal fall. Reapply every 7 to 10 days if bloom is staggered and weather is rainy. Do not use less than 10 gallons of spray volume per acre if aerially applied.	Do not apply more than 19.2 lb KOVERALL (14.4 lb ai) per acre per use season. Do not make last application later than 5 weeks after petal fall. Do not graze livestock in treated area. Do not apply this product with a U-boom device. Minimum retreatment interval: 7 days

10-10-12

1/10/2

Cheminova, Inc.
One Park Drive, Suite 150
Research Triangle Park, NC 27709
www.cheminova.us.com